

AD-A091 124    STANFORD UNIV CA DEPT OF COMPUTER SCIENCE  
LOWER BOUNDS FOR ALGEBRAIC DECISION TREES.(U)  
JUL 80 J M STEELE, A C YAO


F/G 12/1

N00014-76-C-0475  
NL

UNCLASSIFIED    STAN-CS-80-810

1 of 1  
AD-A091 124

END  
DATE  
FILMED  
DTIC



AD A091124

DDC FILE COPY.

Stanford Department of Computer Science  
Report No. STAN-CS-80-810

12  
11 July 1980

14

LEVEL

6

LOWER BOUNDS FOR ALGEBRAIC DECISION TREES

by

10

Michael Steele  
Andrew C. Yao

12  
14

15 N00014-76-C-04753  
✓ NSF-MCS77-05313

Research sponsored by  
National Science Foundation

DTIC  
NOV 3 1980

COMPUTER SCIENCE DEPARTMENT  
Stanford University



DISTRIBUTION STATEMENT A  
Approved for public release;  
Distribution Unlimited

094130  
00 10 20 037

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER STAN-CS-80-810 ✓	2. GOVT ACCESSION NO. AD-A091 184	3. RECIPIENT'S CATALOG NUMBER	
4. TITLE (and Subtitle) Lower Bounds for Algebraic Decision Trees		5. TYPE OF REPORT & PERIOD COVERED technical, July 1980	
7. AUTHOR(s) J. Michael Steele and Andrew C. Yao		6. PERFORMING ORG. REPORT NUMBER STAN-CS-80-810	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Computer Science Stanford University Stanford, California 94305 USA		8. CONTRACT OR GRANT NUMBER(s) ONR N00014-76-C-0475 (NR-042-267) NSF MCS-77-05313-A01	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, Virginia 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
14. MONITORING AGENCY NAME & ADDRESS (if diff. from Controlling Office) ONR Representative - Philip Surra Durand Aeromautic Building, Room 165 Stanford University		12. REPORT DATE July 1980	13. NO. OF PAGES 12
16. DISTRIBUTION STATEMENT (of this report)  Approved for public release; distribution unlimited.		15. SECURITY CLASS. (of this report) Unclassified	
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from report)		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Algebraic Decision trees, Betti Numbers, Connected component, Decision trees, Information theoretic lower bound, Lower bounds.			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  A topological method is given for obtaining lower bounds for the height of algebraic decision trees. The method is applied to the knapsack problem where a $\Omega(n^2)$ bound is obtained for trees with bounded-degree polynomial tests, thus extending the Dobkin-Lipton result for linear trees. Applications to the convex hull problem and the distinct element problem are also indicated. Some open problems are discussed.			

DD FORM 1473  
1 JAN 73  
EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED  
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

## Lower Bounds for Algebraic Decision Trees

J. Michael Steele\* and Andrew C. Yao†

### Abstract

A topological method is given for obtaining lower bounds for the height of algebraic decision trees. The method is applied to the knapsack problem where a  $\Omega(n^2)$  bound is obtained for trees with bounded-degree polynomial tests, thus extending the Dobkin-Lipton result for linear trees. Applications to the convex hull problem and the distinct element problem are also indicated. Some open problems are discussed.

*to George (a friend)*

**Keywords:** Algebraic Decision trees, Betti Numbers, Connected component, Decision trees, Information theoretic lower bound, Lower bounds.

\* Department of Statistics, Stanford University, Stanford, California 94305. The research of this author was supported in part by Office of Naval Research contract N00014-76-C-0475 (NR-042-267).

† Computer Science Department, Stanford University, Stanford, California 94305. The research of this author was supported in part by National Science Foundation grant MCS-77-05313-A01.

## 1. Introduction.

Decision trees are often used to model algorithms for combinatorial and geometrical problems. While motivation for these models rests primarily on their generality and conceptual simplicity, they also have the benefit of offering at present the most promising prospect for proving worst case lower bounds in many problems.

For *linear* decision trees several powerful techniques are known for bounding the tree height from below, e.g. Reingold [9], Dobkin [3], Dobkin and Lipton [4][5], Yao [13], and Yao and Rivest [15].

Much less is known for *general algebraic* decision trees. Beyond the naïve information bound, Rabin's theorem (Rabin [8]) and the convex hull problem (Yao [14]) are apparently the only known results.

The purpose of this article is to provide a general method for establishing lower bounds for the worst case performance of algorithms prescribed by arbitrary algebraic decision trees. Technically this work extends the results of Dobkin and Lipton [4][5], but the tools put to work here provide non-trivial bounds for a large class of previously untouchable problems.

Before giving the detailed computational model it seems worthwhile to mention informally a concrete application.

**Theorem 1.** *Any algebraic decision tree of bounded order which solves the  $n$ -dimensional knapsack problem must have height at least  $\Omega(n^2)$ .*

This result extends the knapsack bounds under the *linear* decision tree model due to Dobkin and Lipton [4] and the  $\Omega(n \log n)$  result of Dobkin [3].

The method used here rests critically on a result from real algebraic geometry due to Milnor [7]. Since the machinery used by Milnor may not be familiar to workers in complexity, we have tried to give an expository of the basic facts necessary for making this work self-contained. The bounds discussed here should prove useful in many related problems.

In the next section we rigorously specify the computational model and outline the lower bound method. The third section exposit Milnor's inequality and gives a

heuristic argument which tries to pinpoint the necessity for the more sophisticated tools.

The fourth section is devoted to applications and in particular to the proof of the result on the knapsack problem (Theorem 1) which was mentioned above.

The final section mentions some open problems and suggest a line of attack which if sufficiently developed might add significantly to the power of the present method.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/ _____	
Availability Codes	
Dist	Avail and/or Special
A	

## 2. Computational Model the General Method.

Let  $W \subseteq R^n$  be any set. A ( $d$ -th order) decision tree  $T$  for testing if  $\bar{x} \in W$  is a ternary tree with each internal node containing a test of the form  $p(x_1, x_2, \dots, x_n) : 0$ , where  $p$  is a polynomial of degree at most  $d$ . Each leaf of  $T$  contains "yes" or a "no" answer. For an input  $\bar{x}$ , the procedure starts at the root and traverses down the tree. At each internal node a branching is made according to the polynomial test at that node and when a leaf is reached the answer to the question "Is  $\bar{x} \in W$ " must be given correctly.

Now let  $C_d(W)$  be the minimum height  $h_T$  for any  $d$ -th order decision tree  $T$  (for the set  $W$ ). Our key objective will be to obtain lower bounds on  $C_d(W)$ , and the bound given here will depend heavily on the topology of  $W$ .

By  $\#W$  we denote the number of (disjoint) connected components of  $W$ . Also for any polynomial  $p(x_1, x_2, \dots, x_n)$  we set  $S_p = \{\bar{x} \mid p(\bar{x}) \neq 0\}$ , and for any integers  $n, m > 0$  we put  $\beta(m, n) = \max\{\#S_p \mid p \text{ is a polynomial of } n \text{ real variables and of degree at most } m\}$ .

The following elementary result provides the skeleton of our method. (To put flesh on the bones will require the bounds on  $\beta$  obtained in the next section.)

**Theorem 2.** Let  $W \subseteq R^n$  be an open set, and let  $T$  be a  $d$ -th order algebraic decision tree for deciding if  $\bar{x} \in W$ . If  $W$  is the disjoint union of  $N$  open sets, then the height  $h_T$  satisfies the inequality

$$2^{h_T} \beta(h_T d, n) \geq N.$$

*Proof.* For each leaf  $\ell$  of  $T$  let  $V_\ell$  be the set of inputs  $\bar{x} \in R^n$  leading to  $\ell$  and let  $I_\ell$  be the set of constraints resulting from the tests. Let  $\mathcal{L}$  be the set of leaves  $\ell$  such that  $I_\ell$  consists only of strict inequalities and such that the answer stored at  $\ell$  is "yes". One should note that each  $V_\ell$  is an open set and  $V_\ell \subseteq W$ .

We now write  $W = \bigcup_{i=1}^N W_i$  where each  $W_i$  is a connected open set and the  $W_i$  are disjoint, and write  $V_\ell = \{\bar{x} : p_{\ell_1}(\bar{x}) < 0, p_{\ell_2}(\bar{x}) < 0, \dots, p_{\ell_s}(\bar{x}) < 0\}$  where each  $p_{\ell_i}$  is a polynomial of degree not greater than  $d$  and where  $s \leq h_T$ . As a consequence of this representation,  $V_\ell \subseteq \{\bar{x} \mid q_\ell(\bar{x}) \neq 0\} = D$



where  $q_\ell(z) = \prod p_{\ell j}(z)$  is a polynomial of degree at most  $h_T d$ . Moreover, each connected component of  $V_\ell$  is contained in at most one component of  $\mathcal{D}$ . Hence,  $V_\ell$  has at most  $\beta(h_T d, n)$  connected components  $V_{\ell 1}, V_{\ell 2}, \dots$ .

Since each leaf of  $T$  is correctly labeled, each  $V_{\ell j}$  has to be completely contained in some  $W_i$ . Since the number of such  $V_{\ell j}$  is at most  $\beta(h_T d, n)$  and there are only  $|\mathcal{L}|$  values of  $\ell$  which lead to "yes" the number of components  $N$  of  $\mathcal{W}$  is bounded by  $|\mathcal{L}| \beta(h_T d, n)$ . Since  $2^{h_T} \geq |\mathcal{L}|$  the theorem follows. ■

### 3. Counting Connected Components.

To use Theorem 2 one needs bounds on  $\beta(m, n)$  and this is apparently no easy matter. Fortunately, there is a bound due to Milnor [7] which is sufficient for some applications:

$$\beta(m, n) \leq (m + 2)(m + 1)^{n-1}. \quad (3.1)$$

The proof of Milnor's inequality rests on the several substantial results from Morse theory and algebraic topology, but it is nevertheless possible to give a heuristic indication of an analogous result.

The only preliminary needed for the argument is Bezout's Theorem which says that any system of  $n$  algebraic equations in  $n$  variable with degree  $d$  has either infinitely many (complex) solutions or at most  $d^n$ . For a classical approach to the proof of Bezout's Theorem one can consult Enriques [6], or, for the case  $n = 2$ , there is a nice proof in Seidenberg [10].

To use Bezout's Theorem we suppose that  $p$  is a real polynomial in  $n$  variables with degree  $m$ , and we note that  $R$  can be chosen so that  $A = \{p > 0\} \cap \{q = R^2 - \sum_{i=1}^n x_i^2 > 0\}$  has as many bounded connected components as  $\{p > 0\}$  has connected components (bounded or unbounded). Since each bounded connected component of  $A$  must contain a local maximum of  $pq$ , the number of bounded components of  $A$  is majorized by the number of zeros of the system  $\nabla pq = 0$ . By Bezout, this number is either infinite, or else bounded by  $(m + 1)^n$ .

This finite bound is for our purposes almost as sharp as Milnor's bound. The real work comes in providing a rigorous perturbation argument which rules out the case when Bezout gives only the trivial infinite bound. That is precisely the case which causes all the trouble and presents this section from being self contained.

We should further remark that a recent exposition of R. Bott [2] provides an intuitive introduction to Milnor [7], where inequality (3.1) is given as Theorem 3. As it happens, the problem of determining  $\beta(m, n)$  is actually very deep and it is intimately connected with Hilbert's 16-th Problem, see Arnold [1].

#### 4. Applications.

We now use Theorem 2 to derive lower bounds. Clearly, the function  $2^z \beta(xd, n)$  is an increasing function of  $x$ . Let  $\alpha(d, n, N)$  be the minimum  $x$  satisfying  $2^z \beta(xd, n) \geq N$ . Theorem 2 immediately yields the following formal bounds:

Any general upper bounds on  $\beta$  can be used to derive lower bounds on  $\alpha$  and hence  $C_d$ . In particular, Milnor's bound (3.1) gives the following result.

**Theorem 3.** For any real  $\epsilon$ ,

$$C_d(W) \geq \min\{\epsilon \log_2 N, \frac{1}{d}(N^{\frac{1-\epsilon}{n}} - 1)\}$$

when  $N = \#W$ .

**Corollary.** If  $\#W = \Omega(n^{(1+\delta)^n})$  for some fixed  $\delta > 0$ , then

$$C_d(W) = \Omega(\log(\#W)).$$

*Proof.* Let  $x = C_d(W)$ . Then  $2^z \beta(xd, n) \geq N$ . Hence by (3.1)

$$2^z (xd + 1)^n \geq N.$$

Either  $2^z \geq N^\epsilon$  or  $(xd + 1)^n \geq N^{1-\epsilon}$ , proving the theorem. ■

The corollary follows by writing  $\#W = n^{n(1+\delta(n))}$  and setting  $\epsilon = \frac{1}{2} \frac{\delta(n)}{1+\delta(n)}$  in the theorem.

Thus, Theorem 3 gives a lower bound nonlinear in  $n$  when  $\#W$  grows at least as fast as  $n^{(1+\delta)^n}$ . This is also necessary since the theorem only gives a lower bound  $O(n)$  when  $\#W = O(n^n)$ .

In the first example given below,  $\#W \approx 2^{\frac{1}{2}n^2}$  thus we have a good lower bound. The other two examples have  $\#W \leq n^n$ , and Theorem 3 does not give

nonlinear bounds. However, Theorem 2 (or, (4.1)) is still true for these later examples, and a better determination in the future may result in an improved bound.

**Example 1. The Knapsack Problem.** Given real numbers  $x_1, x_2, \dots, x_n$ , decide if there exists some subset  $S \subseteq \{1, 2, \dots, n\}$  such that  $\sum_{i \in S} x_i = 1$ .

In this case,  $W = \{(x_1, x_2, \dots, x_n) \mid \prod_S (\sum_{i \in S} x_i - 1) \neq 0\}$ . It was shown in Dobkin's and Lipton (1978)  $\#W \geq 2^{\frac{1}{2}n^2}$ . Thus,  $C_d(W) = \Omega(n^2)$  for any fixed  $d$ . This generalizes the result of Dobkin and Lipton where they showed  $C_1(W) = \Omega(n^2)$ .

**Example 2. Element Distinctness.** Given  $x_1, x_2, \dots, x_n \in \mathcal{R}$ , is there a pair  $i, j$  with  $i \neq j$  and  $x_i = x_j$ ? In this case,

$$W = \{(x_1, x_2, \dots, x_n) \mid \prod_{i \neq j} (x_i - x_j) \neq 0\} \subset \mathcal{R}^n$$

It is easily shown that  $\#W = n!$  since each region  $\{(x_1, x_2, \dots, x_n) \mid x_{\sigma(1)} < x_{\sigma(2)} < \dots < x_{\sigma(n)}\}$  is a maximal connected component of  $W$  for each permutation  $\sigma$ . One therefore has  $C_d(W) \geq \alpha(d, n, n!)$ .

**Example 3. Extreme Points.** Given  $n$  points on the plane does the convex hull formed by them possess  $n$  vertices?

Here  $W$  cannot be expressed by an easy algebraic relation but it is still possible to show  $\#W \geq (n-1)!$ . Obviously,  $W$  is an open set in  $(\mathcal{R}^2)^n$ . For any configuration  $\{x_1, x_2, \dots, x_n\}$  in  $W \subset (\mathcal{R}^2)^n$ , we have a cyclical ordering  $\sigma$  of the points  $\{x_i \mid 1 \leq i \leq n\}$  which is given uniquely by taking the points in cyclical order. Clearly, any of the  $(n-1)!$  cyclical permutations can arise in this way so all that remains is to show that if  $\sigma \neq \sigma'$  then the configurations which give rise to these permutations are in disjoint components of  $W$ .

For each configuration in  $W$  we consider the  $\binom{n}{3}$  element array  $A$  given by  $\Delta(x_i, x_j, x_k)$  where  $\Delta$  is the signed area of the triangle formed by the 3-set  $\{x_i, x_j, x_k\} \subset \{x_1, x_2, \dots, x_n\}$ . If the configuration corresponding to  $\sigma$  is continuously deformed in any way to the configuration for  $\sigma'$  then  $A_\sigma$  is transformed continuously into  $A_{\sigma'}$ . Since  $\sigma$  and  $\sigma'$  differ there is some triple  $\{x_i, x_j, x_k\}$  for which  $\Delta(x_i, x_j, x_k)$  has differing signs in  $A_\sigma$  and  $A_{\sigma'}$ . By the intermediate value

theorem there is therefore some time during the continuous deformation when  $\Delta(x_i, x_j, x_k) = 0$ . This says that  $x_i, x_j, x_k$  are then co-linear and at that point there are at most  $n - 1$  extreme points in the configuration. This proves that any passage from  $\sigma$  to  $\sigma'$  must go out of  $W$ , so  $\sigma$  and  $\sigma'$  correspond to different components.

The main consequence of the preceding bound is that

$$C_d(W) \geq \alpha(d, 2n, (n - 1)!)^2$$

and it was originally hoped that this would be sufficient to prove a conjecture of Yao [14] that any algebraic decision tree of order  $d$  for the extreme point problem must have height  $\Omega(n \log n)$ . The Milnor bound in this case is not sufficiently sharp to obtain the desired bound. We indicate in the next section a bound which would be sufficient.

While these last two examples are disappointing in that they do not give the conjectured non-linear lower-bounds, one should note that since only a yes-no answer is required there is a logical necessity of only 2 terminal leaves. So, the information theoretic bound in these two cases gives only the absurd bound  $\log_2 2$ .

## 5. Open Problems and Directions.

Surely the most interesting and important problems pivot about finding sharper bounds on  $\beta(m, n)$ . It is conceivable that  $\beta(m, n) = 2^{O(m+n)}$  which could imply by Theorem 2 that  $C_d(W) = \Omega(\frac{1}{2}(\log_2 N - m))$ . This bound would yield a  $\Omega(n \log n)$  lower bound in Examples 2 and 3 for fixed  $d$ .

In fact, a somewhat weaker result will suffice for this purpose. Let  $\beta(d, m', n)$  be the maximum of  $\#S_p$  for any  $p$  of the form  $\prod_{i=1}^{m'} p_i(x_1, x_2, \dots, x_n)$ , with each  $p_i$  of degree not greater than  $d$ . Clearly  $\beta(d, m', n) \leq \beta(dm', n)$ . The result one really needs in Examples 2 and 3 is  $\beta(d, m', n) = 2^{O(dm'+n)}$ . Can one prove better bounds on  $\beta(d, m', n)$  than on  $\beta(m, n)$ ? Here we note that it is not hard to see that

$$\beta(1, m', n) \leq \sum_{j=0}^n \binom{m'}{j}, \quad n \leq m' \quad (5.1)$$

since  $\beta(1, m', n)$  just equals the number of regions of  $\mathcal{R}^n$  which can be partitioned by  $m'$  hyperplanes. (This is proved in Steiner (1826) [11] which is in the first volume of Crelle's *J. Reine Ang. Math.* and which is better remembered for containing five fundamental papers of N. H. Abel. For modern treatment of (5.1) see Wetzel [12] and the references given there.)

A more modest approach to the problems suggested by Examples 2 and 3 rest on obtaining bounds for any small values of  $d \geq 2$ . It is known (Yao [14]) that  $C_2(W) = \Omega(n \log n)$  in Example 3, but there are no other known non-linear lower bounds even in the case  $d = 3$ .

## References

- [1] V. Arnold, "Real Algebraic Geometry (th 16-th Hilbert Problem)," *Proc. Symp. Pure Math* 28 (Part 1), 1976, 50-51.
- [2] R. Bott, "Some Recollections from 30 Years Ago," in *Constructive Approaches to Mathematical Models*, (Eds. C. V. Coffman, G. J. Fix), Academic Press, New York, 1979.
- [3] D. Dobkin, "A nonlinear Lower Bound on Linear Search Tree Programs for Solving Knapsack Problems," *J. Comput. System Sci.* 13 (1976), 69-73.
- [4] D. Dobkin and R. J. Lipton, "A Lower Bound of  $\frac{1}{2}n^2$  on Linear Search Programs for the Knapsack Problem," *J. Comput. System Sci.* 16 (1978), 413-417.
- [5] D. Dobkin and R. J. Lipton, "On the Complexity of Varying Sets of Primitives," *J. Comput. and System Sci.* 18 (1980), 86-91.
- [6] F. Enriques, *Lezioni Sulla Teoria Geometrica delle Equazioni e delle Funzioni Algebraiche*, vol. 2, Nicola Zanichelli, Bologna, (1915), 101-104.
- [7] J. Milnor, "On the Betti Numbers of Real Varieties," *Amer. Math. Soc.* 15 (1964), 275-280.
- [8] M. Rabin, "Proving the Simultaneous Positivity of Linear Forms," *J. Comput. System Sci.* 13 (1972), 639-650.
- [9] E. Reingold, "Computing the Maximum and the Median," *Twelfth Annual Symposium on Switching and Automata Theory* (1971), 216-218.
- [10] A. Seidenberg, *Elements of the Theory of Algebraic Curves*, Addison-Wesley, Menlo Park, 1968, p. 44.
- [11] J. Steiner, "Einige Gesetze über die Theilung der Ebene und des Raumes," *J. Reine Angew Math.* (1826), 349-364.
- [12] J. E. Wetzel, "On the Division of the Plane by Lines," *Amer. Math. Monthly* 85 (1978), 647-655.

- [13] A. C. Yao, "On the Complexity of Comparison Problems Using Linear Functions," *Proc. Sixteenth Annual Symposium on the Foundations of Computer Science*, Berkeley, 1975, 85-89.
- [14] A. C. Yao, "A Lower Bound to Finding Convex Hulls," Stanford Computer Science Report STAN-CS-79-733, April 1979 (to appear in *Journal of the ACM*).
- [15] A. C. Yao and R. L. Rivest, "On the Polyhedral Decision Problem," *SIAM J. on Computing* 9 (1980), 343-347.



